VOL. 468 JOURNAL OF CHROMATOGRAPHY MAY 12, 1989

CONTENTS

ADVANCES IN CHROMATOGRAPHY 1988: 25TH INTERNATIONAL SYMPOSIUM, MINNEAPOLIS, MN, AUGUST 29–SEPTEMBER 1, 1989

HONOUR VOLUME ON THE OCCASION OF THE 65TH BIRTHDAY OF ALBERT ZLATKIS

Foreword
by B. S. Middleditch XIII

An introduction and a celebration
by L. S. Ettre XV

Winners of the M.S. Tswett Chromatography medal XVII

1963–1988: Twenty-five years of International Symposia on Advances in Chromatography
by L. S. Ettre (Norwalk, CT, U.S.A.) 1

Chromatography and the discovery process
by C. S. G. Phillips (Oxford, U.K.) 35

Advances in the use of computerized gas chromatography–mass spectrometry and high-performance liquid chromatography with rapid scanning detection for clinical diagnosis
by E. Jellum, O. Thoresen, L. Horn, R. Seip, E. Nilsen, E. A. Kvittingen and O. Stokke (Oslo, Norway) 43

Separation of precolumn *ortho*-phthalaldehyde-derivatized amino acids by capillary zone electrophoresis with normal and micellar solutions in the presence of organic modifiers
by J. Liu, K. A. Cobb and M. Novotny (Bloomington, IN, U.S.A.) 55

Silanediol groups of the silica gel Nucleosil: active sites involved in the chromatographic behaviour of bases
by B. Pfleiderer and E. Bayer (Tübingen, F.R.G.) 67

Stability constants of hydracid–tributyl phosphate complexes in alkane solution: hydrochloric acid and hydrogen cyanide
by K. G. Furton, J. H. Purnell and G. Stedman (Swansea, U.K.) 73

Effective separation of sterol C-24 epimers
by N. Ikekawa (Fukushima, Japan), Y. Fujimoto (Tokyo, Japan) and S. Kadota and T. Kikuchi (Toyama, Japan) 91

Contemporary liquid chromatographic column design
by R. P. W. Scott (Washington, DC, U.S.A. and London, U.K.) 99

Supercritical-fluid chromatography–mass spectrometry of polycyclic aromatic hydrocarbons with a simple capillary direct interface
by S. B. Hawthorne and D. J. Miller (Grand Forks, ND, U.S.A.) 115

Some practical aspects of column design for packed-column supercritical-fluid chromatography
by T. A. Dean and C. F. Poole (Detroit, MI, U.S.A.) 127

Analysis of volatiles in tall oil by gas chromatography, flame-photometric detection, flame-ionization detection and mass spectrometry
by F. Andrawes and T. Chang (Stamford, CT, U.S.A.) and R. Scharrer (Tuxedo, NY, U.S.A.) 145

Gas chromatographic determination of oxo- and hydroxycarboxylic acids in serum and urine of diabetic and normal subjects
by H. M. Liebich, A. Dotzauer and B. Tetschner (Tübingen, F.R.G.) 157

Retention modification of nucleic acid constituents in reversed-phase high-performance liquid chromatography
by R. S. Ramsey (Oak Ridge, TN, U.S.A.), V. W. Chan (Wooster, OH, U.S.A.), B. M. Dittmar (Durham, NC, U.S.A.) and K. H. Row (Knoxville, TN, U.S.A.) 167

Analysis of *in situ* methylated microbial fatty acid constituents by Curie-point pyrolysis–gas chromatography–mass spectrometry
by G. Holzer and T. F. Bourne (Atlanta, GA, U.S.A.) and W. Bertsch (University, AL, U.S.A.) 181

Selectivity of an octadecyl-modified vinyl alcohol copolymer gel for the retention of polar compounds by M. Hirukawa and T. Hanai (Iruma, Japan) 191

Characterisation of products formed by condensations of primary amines with polygodial, and their scope for enantiomer analyses
by C. J. W. Brooks, W. J. Cole and P. A. Brindle (Glasgow, U.K.) 201

Non-radioactive electron-capture detector for gas chromatography
by W. E. Wentworth, T. Limero, C. F. Batten and E. C. M. Chen (Houston, TX, U.S.A.) 215

Analytical-scale high-performance liquid chromatography of omega-3 fatty acid esters derived from fish oils
by J. M. Beebe, P. R. Brown and J. G. Turcotte (Kingston, RI, U.S.A.) 225

Thermodynamic approach to the practical characterization of solvent strength and selectivity of commonly used stationary phases in gas chromatography
by B. R. Kersten, S. K. Poole and C. F. Poole (Detroit, MI, U.S.A.) 235

Changes in retention and polarity accompanying the replacement of hydrogen by fluorine in tetraalkylammonium alkyl- and arylsulfonate salts used as stationary phases in gas chromatography
by R. M. Pomaville and C. F. Poole (Detroit, MI, U.S.A.) 261

Highly sensitive on-line radioassay of high-performance liquid chromatographic effluents
by A. Karmen, G. Malikin and S. Lam (The Bronx, NY, U.S.A.) 279

Simultaneous gas chromatography of volatile and non-volatile carboxylic acids as *tert.*-butyldimethylsilyl derivatives
by K. R. Kim and M. K. Hahn (Seoul, South Korea) and A. Zlatkis, E. C. Horning and B. S. Middleditch (Houston, TX, U.S.A.) 289

Improved flame photometric detector for the analysis of sulfur compounds by gas chromatography
by J. N. Driscoll and A. W. Berger (Newton, MA, U.S.A.) 303

Characterization of coal liquefaction products by gas chromatography–Fourier transform infrared spectroscopy–mass spectrometry
by E. S. Olson and J. W. Diehl (Grand Forks, ND, U.S.A.) 309

Element-selective gas chromatographic detection by atomic plasma emission spectroscopy. Review and developments
by P. C. Uden, Y. Yoo, T. Wang and Z. Cheng (Amherst, MA, U.S.A.) 319

Isocratic ion-pair high-performance liquid chromatographic methods for the determination of uroporphyrin and coproporphyrin type II and IV isomers in human urine
by K. Jacob, E. Egeler, D. Neumeier and M. Knedel (Munich, F.R.G.) 329

Method for analysis of dilute vapours in flue gases and working atmospheres
by E. D. Morgan and N. Bradley (Keele, U.K.) 339

Determination of the total amount sulfur in petroleum fractions by capillary gas chromatography in combination with cold trapping and a total sulfur analyzer
by N. J. Sung and S. J. Johnson (Bartlesville, OK, U.S.A.) 345

Confirming chiral high-performance liquid chromatographic separations with stereospecific enzymes
by S. Lam, G. Malikin, M. Murphy, L. Freundlich and A. Karmen (The Bronx, NY, U.S.A.) 359

Wall-coated adsorbent open-tubular fused-silica columns for the separation of complex pyrolysis products
by M. Xiang (Beijing, China) and F. Bruner (Urbino, Italy) 365

Dispersion and selectivity indices in gas chromatography. IV. Chlorinated aromatic compounds
by M. B. Evans (Hatfield, U.K.) and J. K. haken (Kensington, Australia) 373

Microdetermination of 11-dehydrothromboxane B_2 in human urine by gas chromatography–selected-ion monitoring
by K. Watanabe, K. Yamashita and M. Ishibashi (Tokyo, Japan), Y. Hayashi and S. Yamamoto (Tokushima, Japan) and H. Miyazaki (Tokyo, Japan) 383

Author Index 395

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)